

Б Р Б Ч С Б Р Э Р 2 Р 4 С В Р Р Б У С Р S П S
ЕРЕВАНСКИЙ ФИЗИЧЕСКИЙ ИНСТИТУТ

ЕФМ-ТФ-ИИ(71)

V.A. Khoze, I.A. Nagorskaya and A.N. Zaslavsky

ELECTROMAGNETIC DECAYS OF $X^0(960)$ AND
E(1420) MESONS

А Р У С



Б Р Б Ч С Б Р

1971

Е Р Е В А Н

Yerevan Physical Institute

EQM-10-II(71)

ELECTROMAGNETIC DECAYS OF $\chi^0(960)$ AND $E(1420)$ MESONS

V.A. Khoze, I.A. Nagorskaya and A.N. Zaslavsky

YEREVAN

1971

ABSTRACT

The electromagnetic decays of the $\chi^0(960)$ and $E(1420)$ mesons have been studied. The decays with the Dalitz pair production have been analyzed in detail.

Introduction

There are two main candidates, the $X^0(960)$ and the $E(1420)$ mesons, at present for the ninth pseudoscalar meson [1,2]. The spin-parity of the $X^0(960)$ meson has not been yet firmly established, and of the two possible hypotheses $J^P(X^0) = 0^-$ and 2^- , the latter is even more preferable than the former [2]. On the other hand, the alternative hypothesis for the $E(1420)$ meson spin-parity is $J^P(E) = 1^+$. In the paper [3] an estimation for the $E \rightarrow 2\gamma$ decay probability has been obtained assuming $E-\eta$ mixing. The importance of the experimental study of this process is emphasized as the detection of the $E \rightarrow 2\gamma$ decay would unambiguously solve the problem of the $E(1420)$ meson spin-parity in favour of the 0^- hypothesis and completely exclude the other possible 1^+ hypothesis.

The interest for the experimental investigation of the $X^0 \rightarrow 2\gamma$ [4,5] has been recently increased. The experimental study of the $E(1420)$ radiative decays may be appeared very interesting, since, according to the available estimations [3], the branching ratios for these decays are not small.

In this paper we shall mainly consider the rare radiative decays of the X^0 and E mesons with Dalitz pair production. In cases when it makes sense, the possibility for determining the spin-parity of the X^0 and E mesons by their radiative decay will be emphasized.

I. The radiative decays of the $X^0(960)$ meson

I. The $X^0 \rightarrow \rho^0 e^+ e^-$ decay.

For the $J^P(X^0) = 0^-$ hypothesis the formula for the width of the $X^0 \rightarrow \rho^0 \gamma$ decay is well known:

$$\Gamma(X^0 \rightarrow \rho\gamma) = |g_{X\rho\gamma}|^2 \frac{(M_X^2 - M_\rho^2)^3}{32\pi M_X^3}, \quad (1)$$

where $g_{X\rho\gamma}$ is determined by the matrix element of the $X^0 \rightarrow \rho\gamma$ decay:

$$M_{0-}(X^0 \rightarrow \rho\gamma) = \frac{g_{X\rho\gamma}}{4} \varphi(Q) \tilde{F}_{\mu\nu}(\Delta) p_{\mu\nu}(q) \quad (2)$$

Assuming that the form factor $g_{X\rho\gamma}(\Delta^2)$, where $\Delta^2 = (p_e + p_{e^-})^2$, is a constant [6,7] at small energy separation (~ 200 MeV), it will be easy to calculate the conversion coefficient $K = \Gamma(X^0 \rightarrow \rho e^+ e^-) / \Gamma(X^0 \rightarrow \rho\gamma)$ and the Dalitz plot distribution for the $X^0 \rightarrow \rho e^+ e^-$ decay with an accuracy to $\Delta^2/M_X^2 < 1/25$ ($4m_e^2 \leq \Delta^2 \leq (M_X - M_\rho)^2$):

$$K = \frac{\Gamma(X^0 \rightarrow \rho e^+ e^-)}{\Gamma(X^0 \rightarrow \rho\gamma)} = \frac{2}{3\pi} \left(2 \ln \frac{M_X - M_\rho}{m_e} - 3,4 \right) \approx \frac{1}{150} \quad (3)$$

$$dW = \frac{2M_X}{(2\pi)^2} |g_{X\rho\gamma}(\Delta^2)|^2 \left\{ (\Delta^2 + 2m_e^2) \left[(E_1 + E_2)^2 - \Delta^2 \right] + \Delta^2 \left(\frac{\Delta^2}{2} - 2E_1 E_2 \right) \right\} \frac{dE_1 dE_2}{\Delta^4} \quad (4)$$

Here E_1 (E_2) is the electron (positron) energy in the X^0 meson rest frame and $\Delta^2 = 2M_X(E_1 + E_2) + M_\rho^2 - M_X^2$.

As the energy separation is small, the hypothesis 0^- cannot be distinguished from 2^- for the $X^0(960)$ spin-parity by means of the magnitude of the conversion coefficient K and the Dalitz plot for the $X^0 \rightarrow \rho e^+ e^-$ decay [8]. One can show that the conversion coefficient and the Dalitz plot distribution for the $X^0 \rightarrow \rho e^+ e^-$ in the case of $J^P(X^0) = 2^-$ coincide within an accuracy up to the terms $\Delta^2/M_X^2 < 1/25$ with the formulae (3) and (4) obtained on the assumption that the X^0 meson is pseudoscalar.

However, the study of the correlation between the relative momentum of pions from the ρ^0 meson decay and the momenta of $e^+ e^-$ pair in the ρ^0 meson rest frame allows to distinguish between hypothesis 0^- and 2^- for $X^0(960)$.

For the alternative $J^P(X^0) = 0^-$ the dependence of the distribution of the decay pions on the angle of emission is sharply expressed

$$W(\theta, \varphi, \gamma) = \left[\frac{\Delta^2}{2} - 2 \sin^2 \theta \sin^2 \varphi \bar{p}_+^2 \right] \sin^2 \gamma, \quad (5)$$

where θ is the angle between the directions of the positron momentum \vec{p}_+ and the X^0 meson momentum \vec{K} in ρ^0 rest frame, γ is the angle between the directions of the X^0 and the π^+ in the same frame, φ is the angle between the planes formed by the vectors \vec{k}, \vec{p} and \vec{k}, \vec{q} .

A good agreement of the experimental data on the $X^0 \rightarrow \rho \gamma$ decay with the $J^P(X^0) = 2^-$ hypothesis is achieved with the help of a simple matrix element [1,2]

$$M_{2^-}(X^0 \rightarrow \rho \gamma) = g_1 T_{\mu\nu}(Q) F_{\mu\lambda}(\Delta) \beta_{\lambda\nu}(q) \quad (6)$$

The distribution $W(\theta, \varphi, \gamma)$ calculated using this matrix element, has the form

$$W(\theta, \varphi, \gamma) = \frac{1}{2} (\Delta^2 - 2\bar{p}^2 \sin^2 \theta) + \frac{1}{6} \frac{2M_\rho^2 - M_X^2}{M_X^2} \sin^2 \gamma \left[\frac{\Delta^2}{2} - 2\bar{p}^2 \sin^2 \theta \sin^2 \varphi \right] - \Delta^2 \frac{M_X^2 + M_\rho^2}{2M_X^2} \left[\left(\frac{M_X^2 - M_\rho^2}{2M_X^2} - \frac{|F|}{M_\rho} \right) \cos \gamma + \frac{2\rho \cdot \Delta^0 |F|}{|\Delta| M_\rho} \cos \psi \right]$$

$\cos \psi = \cos \theta \cos \gamma + \sin \theta \sin \gamma \cos \varphi$ and depends weakly on the angles φ and γ . Thus, the study of the $X^0 \rightarrow \rho e^+ e^-$ decay in case of sufficient statistics allows to distinguish between the hypotheses 0^- and 2^- for the spin-parity of X^0 (960) meson.

2. The $X^0 \rightarrow \rho e^+ e^-$, $X^0 \rightarrow e^+ e^-$ and $X^0 \rightarrow \pi^+ \pi^- \pi^0$ decays

The $X^0 \rightarrow \gamma \gamma$ decay has been experimentally studied in the reactions $\pi^+ p \rightarrow n X^0$ [4] and $\pi^+ d \rightarrow p(p) + 2\gamma$ [5]. The averaged branching ratio for this decay is $\Gamma(X^0 \rightarrow 2\gamma) = 9,9^{+0,4}_{-3,9} \%$. The conversion coefficient $k_2 = \Gamma(X^0 \rightarrow e^+ e^-) / \Gamma(X^0 \rightarrow \gamma \gamma)$ estimated by the Dalitz formula [6] on the assumption that X^0 meson is pseudo-

scalar, is equal to about 1/60. In this case, however, because the energy separation is great, the form factor $g_{\chi\rho\gamma}(\Delta^2)$ must be taken into account. We have estimated the conversion coefficient in the vector dominance model. For the vertices $g_{\chi\rho\gamma}$, $g_{\chi\omega\gamma}$,

$g_{\chi\eta\gamma}$ we have used the relations of $SU(3)$ and $SU_w(6)$ symmetries. The ratio $\Gamma(\chi^0 \rightarrow \chi e^+ e^-) / \Gamma(\chi^0 \rightarrow \rho\gamma)$ has appeared to be equal to $16 \cdot 10^{-3}$

while the relative decay probability $\Gamma(\chi^0 \rightarrow \chi e^+ e^-) / \Gamma(\chi^0 \rightarrow \text{all}) \approx 0.5 \cdot 10^{-3}$.

For the two possible hypotheses $J^P(\chi^0) = 0^-$ and 2^- , the $\chi^0 \rightarrow e^+ e^-$ decay is suppressed by a factor of $(m_e/m_\chi)^2 \approx 10^{-6}$ since $CP(\chi^0) = -1$ [9].

If $J^P(\chi^0) = 2^-$, then the strong decay $\chi^0 \rightarrow 2\pi$ is suppressed because of the centrifugal barrier, and the relative contribution of the radiative decay modes must be great [2]. This qualitative remark is in agreement with the experimental values: $\Gamma(\chi^0 \rightarrow \rho\gamma) / \Gamma(\chi^0 \rightarrow \text{all}) \sim 30\%$, $\Gamma(\chi^0 \rightarrow \gamma\gamma) / \Gamma(\chi^0 \rightarrow \text{all}) \sim 10\%$. In this case the probability of the electromagnetic decay $\chi^0 \rightarrow \pi^+ \pi^- \pi^0$ may also appear to be not small (about some per cent). The experimental observation and the study of the Dalitz plot of the electromagnetic decay $\chi^0 \rightarrow \pi^+ \pi^- \pi^0$ would help to clear up the question concerning the χ^0 meson spin as it was succeeded in obtaining some arguments in favour of pseudoscalar η -meson by the help of the decay $\eta \rightarrow \pi^+ \pi^- \pi^0$.

II. The radiative decays of $\mathbb{E}(1420)$ meson

I. The decays $\mathbb{E} \rightarrow 2\gamma$, $\mathbb{E} \rightarrow \rho\gamma$

As to the existing alternative spin value for $\mathbb{E}(1420)$, it is very important to study the radiative decays $\mathbb{E} \rightarrow 2\gamma$, $\mathbb{E} \rightarrow \rho\gamma$, etc.

The probability of the decay $\mathbb{E} \rightarrow 2\gamma$, estimated on the assumption about the \mathbb{E} - η mixing with mixing angle equal to -6.5° [3], is great. $\Gamma(\mathbb{E} \rightarrow 2\gamma) \sim (300 - 1000) \text{ KeV}$ and $\Gamma(\mathbb{E} \rightarrow \rho\gamma) \sim 10 \text{ MeV}$.

These circumstances are in favour of the experimental search for E-meson radiative decays. The detection of the decay $E \rightarrow 2\gamma$ would definitively prove that E(I420) is pseudoscalar and completely exclude the other possible value $J^P(E) = 1^+$

The detection of the decay $E \rightarrow \rho\gamma$ and the investigation of its distribution with respect to the angle θ between the relative momentum of the pions and the γ -quantum momentum in ρ meson rest frame is of interest. If $J^P(E) = 0^-$, then $W(\theta) \sim \sin^2\theta$. The deviation from the distribution $\sin^2\theta$ is incompatible with the pseudoscalarity of E(I420) meson. For the hypothesis $J^P(E) = 1^+$, the amplitude of the transition M1 leads to the distribution $W(\theta) \sim 1 + \cos^2\theta$ while the simple relativistic matrix element $T = g b_\mu(p) \tilde{F}_{\nu\lambda}(k) p_\nu(q)$ (b_μ and p_ν are the polarization of E and ρ mesons, respectively) gives

$$W(\theta) \sim 1 + \cos^2\theta - \frac{(M_E^2 - M_\rho^2)}{M_E^2} \sin^2\theta \sim \cos^2\theta.$$

2. The decays $E \rightarrow \rho e^+ e^-$, $E \rightarrow \gamma e^+ e^-$ and $E \rightarrow e^+ e^-$

If E(I420) is pseudoscalar, then the conversion coefficient $K_2 = \Gamma(E \rightarrow \rho e^+ e^-) / \Gamma(E \rightarrow \rho\gamma)$ estimated by a formula analogous to (3), is equal to $1/15$. Any deviation of the distribution $W(\theta, \phi, \gamma)$ for the decay $E \rightarrow \rho e^+ e^-$ from the distribution given by the formula (5) shows an evidence for $J^P(E) = 1^+$.

For the decay $E \rightarrow \gamma e^+ e^-$, the Dalitz formula [6] leads to $K_3 = \Gamma(E \rightarrow \gamma e^+ e^-) / \Gamma(E \rightarrow \gamma\gamma) = 1/45$ while the vector dominance model gives $K_4 = \Gamma(E \rightarrow \gamma e^+ e^-) / \Gamma(E \rightarrow \rho\gamma) = 3.5 \cdot 10^{-4}$.

For the alternative $J^P(E) = 0^-$ the decay $E \rightarrow e^+ e^-$ is suppressed by a factor $(m_e/M_E)^2 \approx 10^{-7}$ due to the helicity conservation. The detection of the decay $E \rightarrow e^+ e^-$, therefore, would show an evidence for $J^P(E) = 1^+$.

III. $M^0(953)$ meson

An evidence [10] for the existence of a new $M^0(953)$ resonance decaying into $2\pi^+\pi^-$ and $\pi^+\pi^0$ has been recently obtained in the reaction $K^-\rho \rightarrow K^-\rho M^0$. The difference of $M^0(953)$ from $X^0(960)$ is that the $\pi^+\pi^-$ effective mass distribution in the $M^0 \rightarrow \pi^+\pi^-$ is isotropic. The possible existence of the $M^0(953)$ requires careful analysis of the final $\pi\pi\gamma$ state [1].

The suggested explanation of this fact [1] accepts that $T_{J^P}^{G(M)} = 0^+ 1^+$ and the decay $M^0 \rightarrow \rho\gamma$ is suppressed in vector dominance model. This explanation is incorrect since in this case the vector dominance model does not limit the number of the spin amplitudes.

The most essential is now the questions about the isospin and the C -parity of the new resonance. Since the decay $M^0 \rightarrow \pi^+\pi^-$ was observed, then $G(M) = 1$ with a high probability. As the pions effective mass distribution is isotropic, then it will be more natural to make assumption that $C(M) = -1$ and $T(M) = 1$. In this case the detection of the decay $M^0 \rightarrow 2\pi^0\gamma$ and the search for $M(953)$ in the reaction $K^-\rho \rightarrow K^0\rho M^0$ are important.

The authors are grateful to S.B. Gerasimov, K.A. Ispirian, S.H. Matinian, B.N. Valuev, V.F. Vishnevsky and especially V.I. Ogievetsky for helpful discussions.


References

1. Particle Data Group, Rev. Mod. Phys. 43, April (1971).
2. V.Ogievetsky, W.Tybor and A.Zaslavsky, Yadernaya Fiz. 9, 852 (1969); Phys. Letters 35B, 69 (1971).
3. A.N.Zaslavsky and W.Tybor, Acta Phys. Polonica 35, 777 (1969); Yadernaya Fiz. 12, 376 (1970).
4. D.Bollini, A.Buhler-Broglin, P.Dalpiaz, T.Massam, F.Navach and Zichichi, Nuovo Cimento 58A, 289 (1968).
5. J.Bensinger, A.R.Erwin, M.A.Thompson and W.D.Walker, Phys. Letters 33B, 505 (1970).
6. R.H.Dalitz, Proc. Phys. Soc. A64, 667 (1951).
7. B.N.Valuev, Preprint JINR (Dubna) P-2823 (1966)
8. V.A.Khoze and A.N.Zaslavsky, Zh. Exper. Teor. Fiz. Pis'ma 13, 706, (1971).
9. V.A.Khoze, Zh. Exper. Teor. Fiz. Pis'ma, 12, 542 (1970).
10. M.Aguilar-Benitez, D.Bassano, R.L.Eisner, J.B.Kinson, D.Pandoulas, N.P.Samios, and V.E.Barnes, Phys. Rev. Letters 25, 1635 (1970).
11. J.L.Rosner and E.W.Colglazier, Preprint Minnesota, Caltech - Collaboration January (1971).

Translated by M.Levonian

Received by Publishing Department

on November 4, 1971



Рукопись поступила 2-го ноября 1971г.

Заказ 681

T-03695

Тираж 300

Множительно-копировальная станция Ереванского физического
института, Ереван 36, Маркаряна 2